

# Lower order eigenvalues of the poly-Laplacian with any order on spherical domains <sup>\*</sup>

Guangyue Huang, Bingqing Ma <sup>†</sup>

Department of Mathematics, Henan Normal University, Xinxiang 453007, Henan  
People's Republic of China

**Abstract.** We consider the lower order eigenvalues of poly-Laplacian with any order on spherical domains. We obtain universal inequalities for them and show that our results are optimal.

**Keywords:** eigenvalue; poly-Laplacian.

**Mathematics Subject Classification:** Primary 35P15; Secondary 53C20.

## 1 Introduction

Let  $\Omega$  be a connected bounded domain in an  $n$ -dimensional complete Riemannian manifold  $M$ . In this paper, we consider the Dirichlet eigenvalue problem of the poly-Laplacian with order  $p$ :

$$\begin{cases} (-\Delta)^p u = \lambda u & \text{in } \Omega, \\ u = \frac{\partial u}{\partial \nu} = \cdots = \frac{\partial^{p-1} u}{\partial \nu^{p-1}} = 0 & \text{on } \partial\Omega, \end{cases} \quad (1.1)$$

where  $\Delta$  is the Laplacian in  $M$  and  $\nu$  denotes the outward unit normal vector field of  $\partial\Omega$ . Let  $0 < \lambda_1 \leq \lambda_2 \leq \lambda_3 \leq \cdots \rightarrow +\infty$  denote the successive eigenvalues for (1.1), where each eigenvalue is repeated according to its multiplicity.

When  $p = 1$ , the eigenvalue problem (1.1) is called a *fixed membrane problem*. For  $M = \mathbb{R}^2$  and  $p = 1$ , Payne-Pólya-Weinberger [10] proved

$$\lambda_2 + \lambda_3 \leq 6\lambda_1.$$

In 1993, for general dimensions  $n \geq 2$ , Ashbaugh and Benguria [2] proved

$$\sum_{i=1}^n (\lambda_{i+1} - \lambda_1) \leq 4\lambda_1.$$

Recently, the inequalities of eigenvalues of the fixed membrane problem have been generalized to some Riemannian manifolds. For the related research and improvement

---

<sup>\*</sup>This research is supported by Project of Henan Provincial department of Sciences and Technology (No. 092300410143), and NSF of Henan Provincial Education department (No. 2009A110010).

<sup>†</sup>The corresponding author. Email: bqma@henannu.edu.cn

in this direction, see [3, 4, 9, 11] and the references therein. In particular, Sun-Cheng-Yang [11] proved that when  $M$  is an  $n$ -dimensional unit sphere  $\mathbb{S}^n(1)$ ,

$$\sum_{i=1}^n (\lambda_{i+1} - \lambda_1) \leq 4\lambda_1 + n^2. \quad (1.2)$$

When  $p = 2$ , the eigenvalue problem (1.1) is called a *clamped plate problem*. For  $M = \mathbb{R}^n$  and  $p \geq 2$ , Cheng-Ichikawa-Mametsuka proved in [8] that

$$\sum_{i=1}^n (\lambda_{i+1} - \lambda_1) \leq 4p(2p-1)\lambda_1, \quad (1.3)$$

$$\sum_{i=1}^n (\lambda_{i+1}^{\frac{1}{p}} - \lambda_1^{\frac{1}{p}})^{p-1} \leq (2p)^{p-1} \lambda_1^{\frac{p-1}{p}}. \quad (1.4)$$

Inequalities (1.3) and (1.4) include two universal inequalities of the clamped plate problem announced by Ashbauth in [1]. When  $M$  is a general complete Riemannian manifold, for  $p = 2$ , Cheng-Huang-Wei [6] obtained

$$\sum_{i=1}^n (\lambda_{i+1} - \lambda_1)^{\frac{1}{2}} \leq (4\lambda_1^{\frac{1}{2}} + n^2 H_0^2)^{\frac{1}{2}} \{(2n+4)\lambda_1^{\frac{1}{2}} + n^2 H_0^2\}^{\frac{1}{2}}, \quad (1.5)$$

where  $H_0^2$  is a nonnegative constant which depends only on  $M$  and  $\Omega$ . For  $M = \mathbb{S}^n(1)$ , we have  $H_0^2 = 1$  such that (1.5) becomes the following inequality:

$$\sum_{i=1}^n (\lambda_{i+1} - \lambda_1)^{\frac{1}{2}} \leq (4\lambda_1^{\frac{1}{2}} + n^2)^{\frac{1}{2}} \{(2n+4)\lambda_1^{\frac{1}{2}} + n^2\}^{\frac{1}{2}}. \quad (1.6)$$

We remark that when  $\Omega = \mathbb{S}^n(1)$ , it holds that  $\lambda_1 = 0$  and  $\lambda_2 = \dots = \lambda_{n+1} = n^2$ . Therefore, the inequality (1.6) becomes equality. Hence, for  $M = \mathbb{S}^n(1)$ , the inequality (1.6) is optimal.

In the present article, we consider the eigenvalue problem (1.1) with any  $p$  when  $M$  is a unit sphere  $\mathbb{S}^n(1)$ . we obtain the following result:

**Theorem.** *Let  $\Omega$  be a bounded domain in an  $n$ -dimensional unit sphere  $\mathbb{S}^n(1)$ . Let  $\lambda_i$  be the  $i$ -th eigenvalue of the eigenvalue problem (1.1). Then we have*

$$\begin{aligned} \sum_{i=1}^n (\lambda_{i+1} - \lambda_1)^{\frac{1}{2}} &\leq \left\{ \left( \lambda_1^{\frac{1}{p}} + n \right)^p - \lambda_1 + 4[2^p - (p+1)]\lambda_1^{\frac{1}{p}} \left( \lambda_1^{\frac{1}{p}} + n \right)^{p-2} \right\}^{\frac{1}{2}} \\ &\quad \times \left\{ 4\lambda_1^{\frac{1}{p}} + n^2 \right\}^{\frac{1}{2}}. \end{aligned} \quad (1.7)$$

**Remark 1.** *For  $p = 2$ , the inequality (1.7) becomes the optimal inequality (1.6).*

**Remark 2.** *For the unit sphere  $\mathbb{S}^n(1)$ , by taking  $\Omega = \mathbb{S}^n(1)$ , we know  $\lambda_1 = 0$  and  $\lambda_2 = \dots = \lambda_{n+1} = n^2$ . Hence, the inequality (1.7) becomes equality. Therefore, our result is optimal.*

**Acknowledgement.** *The authors thank Professor Qing-Ming Cheng for his helpful discussion and support.*

## 2 Proof of Theorem

Let  $u_i$  be the orthonormal eigenfunction corresponding to eigenvalue  $\lambda_i$ , that is,

$$\begin{cases} (-\Delta)^p u_i = \lambda_i u_i & \text{in } \Omega, \\ u_i = \frac{\partial u_i}{\partial \nu} = \cdots = \frac{\partial^{p-1} u_i}{\partial \nu^{p-1}} = 0 & \text{on } \partial\Omega, \\ \int_{\Omega} u_i u_j = \delta_{ij}. \end{cases}$$

Let  $x^1, x^2, \dots, x^{n+1}$  be the standard Euclidean coordinate functions of  $\mathbb{R}^{n+1}$ , then

$$\mathbb{S}^n(1) = \left\{ (x^1, x^2, \dots, x^{n+1}) \in \mathbb{R}^{n+1} ; \sum_{i=1}^{n+1} (x^i)^2 = 1 \right\}.$$

It is well known that

$$\Delta x^i = -n x^i, \quad i = 1, 2, \dots, n+1.$$

Assume that  $B$  is an  $(n+1) \times (n+1)$ -matrix defined by  $B = (b_{ij})$ , where

$$b_{ij} = \int_{\Omega} x^i u_1 u_{j+1}.$$

Using the orthogonalization of Gram and Schmidt, we know that there exist an upper triangle matrix  $R = (r_{ij})$  and an orthogonal matrix  $Q = (q_{ij})$  such that  $R = QB$ , that is,

$$r_{ij} = \sum_{k=1}^{n+1} q_{ik} b_{kj} = \sum_{k=1}^{n+1} \int_{\Omega} q_{ik} x^k u_1 u_j = 0, \quad 2 \leq j \leq i \leq n+1.$$

Defining  $h_i = \sum_{k=1}^{n+1} q_{ik} x^k$ , one gets

$$\int_{\Omega} h_i u_1 u_j = \sum_{k=1}^{n+1} \int_{\Omega} q_{ik} x^k u_1 u_j = 0, \quad 2 \leq j \leq i \leq n+1.$$

Setting

$$\varphi_i = h_i u_1 - u_1 \int_{\Omega} h_i u_1^2.$$

Then

$$\int_{\Omega} \varphi_i u_j = 0, \quad \text{for any } j \leq i.$$

It follows from Rayleigh-Ritz inequality that

$$\lambda_{i+1} \leq \frac{\int_{\Omega} \varphi_i (-\Delta)^p \varphi_i}{\|\varphi_i\|^2}, \quad (2.1)$$

where  $\|f\|^2 = \int_{\Omega} |f|^2$ . By a direct calculation, we derive at

$$\begin{aligned}
\int_{\Omega} \varphi_i (-\Delta)^p \varphi_i &= \int_{\Omega} \varphi_i (-\Delta)^p (h_i u_1) \\
&= \int_{\Omega} \varphi_i \{ [(-\Delta)^p (h_i u_1) - h_i (-\Delta)^p u_1] + \lambda_1 h_i u_1 \} \\
&= \lambda_1 \|\varphi_i\|^2 + \int_{\Omega} \varphi_i [(-\Delta)^p (h_i u_1) - h_i (-\Delta)^p u_1] \\
&= \lambda_1 \|\varphi_i\|^2 + \int_{\Omega} h_i u_1 [(-\Delta)^p (h_i u_1) - h_i (-\Delta)^p u_1] \\
&\quad - \int_{\Omega} h_i u_1^2 \int_{\Omega} u_1 [(-\Delta)^p (h_i u_1) - h_i (-\Delta)^p u_1] \\
&= \lambda_1 \|\varphi_i\|^2 + \int_{\Omega} h_i u_1 [(-\Delta)^p (h_i u_1) - h_i (-\Delta)^p u_1].
\end{aligned} \tag{2.2}$$

Defining

$$\nabla^r = \begin{cases} \Delta^{r/2} & \text{when } r \text{ is even,} \\ \nabla(\Delta^{(r-1)/2}) & \text{when } r \text{ is odd.} \end{cases}$$

Then (2.2) can be written as

$$\int_{\Omega} \varphi_i (-\Delta)^p \varphi_i = \lambda_1 \|\varphi_i\|^2 + \|\nabla^p (h_i u_1)\|^2 - \lambda_1 \|h_i u_1\|^2. \tag{2.3}$$

Putting (2.3) into (2.1) yields

$$(\lambda_{i+1} - \lambda_1) \|\varphi_i\|^2 \leq \|\nabla^p (h_i u_1)\|^2 - \lambda_1 \|h_i u_1\|^2. \tag{2.4}$$

One gets from integration by parts that

$$\begin{aligned}
\int_{\Omega} u_1 h_i \langle \nabla h_i, \nabla u_1 \rangle &= \frac{1}{4} \int_{\Omega} \langle \nabla (h_i^2), \nabla (u_1^2) \rangle = -\frac{1}{4} \int_{\Omega} u_1^2 \Delta (h_i^2) \\
&= -\frac{1}{2} \int_{\Omega} u_1^2 h_i \Delta h_i - \frac{1}{2} \int_{\Omega} u_1^2 |\nabla h_i|^2.
\end{aligned}$$

Hence,

$$\begin{aligned}
& \int_{\Omega} \varphi_i \left( \langle \nabla h_i, \nabla u_1 \rangle + \frac{1}{2} u_1 \Delta h_i \right) \\
&= \int_{\Omega} u_1 h_i \langle \nabla h_i, \nabla u_1 \rangle + \frac{1}{2} \int_{\Omega} u_1^2 h_i \Delta h_i - \int_{\Omega} h_i u_1^2 \left( \int_{\Omega} u_1 \langle \nabla h_i, \nabla u_1 \rangle + \frac{1}{2} \int_{\Omega} u_1^2 \Delta h_i \right) \\
&= \int_{\Omega} u_1 h_i \langle \nabla h_i, \nabla u_1 \rangle + \frac{1}{2} \int_{\Omega} u_1^2 h_i \Delta h_i \\
&= -\frac{1}{2} \int_{\Omega} u_1^2 |\nabla h_i|^2 \\
&= -\frac{1}{2} \|u_1 \nabla h_i\|^2.
\end{aligned} \tag{2.5}$$

By virtue of (2.4) and (2.5), it is easy to see

$$\begin{aligned}
(\lambda_{i+1} - \lambda_1)^{\frac{1}{2}} \|u_1 \nabla h_i\|^2 &= -2(\lambda_{i+1} - \lambda_1)^{\frac{1}{2}} \int_{\Omega} \varphi_i \left( \langle \nabla h_i, \nabla u_1 \rangle + \frac{1}{2} u_1 \Delta h_i \right) \\
&\leq \delta (\lambda_{i+1} - \lambda_1) \|\varphi_i\|^2 + \frac{1}{\delta} \left\| \langle \nabla h_i, \nabla u_1 \rangle + \frac{1}{2} u_1 \Delta h_i \right\|^2 \\
&\leq \delta \{ \|\nabla^p(h_i u_1)\|^2 - \lambda_1 \|h_i u_1\|^2 \} + \frac{1}{\delta} \left\| \langle \nabla h_i, \nabla u_1 \rangle + \frac{1}{2} u_1 \Delta h_i \right\|^2,
\end{aligned} \tag{2.6}$$

where  $\delta$  is a positive constant. Summing over  $i$  from 1 to  $n+1$  for (2.6), one finds that

$$\begin{aligned}
\sum_{i=1}^{n+1} (\lambda_{i+1} - \lambda_1)^{\frac{1}{2}} \|u_1 \nabla h_i\|^2 &\leq \delta \sum_{i=1}^{n+1} \{ \|\nabla^p(h_i u_1)\|^2 - \lambda_1 \|h_i u_1\|^2 \} \\
&\quad + \frac{1}{\delta} \sum_{i=1}^{n+1} \left\| \langle \nabla h_i, \nabla u_1 \rangle + \frac{1}{2} u_1 \Delta h_i \right\|^2 \\
&= \delta \sum_{i=1}^{n+1} \{ \|\nabla^p(x_i u_1)\|^2 - \lambda_1 \|x_i u_1\|^2 \} \\
&\quad + \frac{1}{\delta} \sum_{i=1}^{n+1} \left\| \langle \nabla x_i, \nabla u_1 \rangle + \frac{1}{2} u_1 \Delta x_i \right\|^2 \\
&= \delta \sum_{i=1}^{n+1} \int_{\Omega} u_1 x_i \{ (-\Delta)^p(u_1 x_i) - x_i (-\Delta)^p u_1 \} \\
&\quad + \frac{1}{\delta} \sum_{i=1}^{n+1} \left\| \langle \nabla x_i, \nabla u_1 \rangle + \frac{1}{2} u_1 \Delta x_i \right\|^2.
\end{aligned} \tag{2.7}$$

Making use of the same method as proof of Lemma 1 in [5], it is easy to prove

$$\int_{\Omega} |\nabla u_1|^2 \leq \lambda_1^{\frac{1}{p}}.$$

Thus,

$$\begin{aligned}
\sum_{i=1}^{n+1} \left\| \langle \nabla x_i, \nabla u_1 \rangle + \frac{1}{2} u_1 \Delta x_i \right\|^2 &= \sum_{i=1}^{n+1} \int_{\Omega} \left( \langle \nabla x_i, \nabla u_i \rangle + \frac{1}{2} u_1 \Delta x_i \right)^2 \\
&= \sum_{i=1}^{n+1} \int_{\Omega} \left( \frac{1}{4} u_1^2 (\Delta x_i)^2 + \langle \nabla x_i, \nabla u_1 \rangle^2 + \frac{1}{2} \Delta x_i \langle \nabla x_i, \nabla (u_1^2) \rangle \right) \\
&= \frac{n^2}{4} + \int_{\Omega} |\nabla u_1|^2 \\
&\leq \frac{n^2}{4} + \lambda_1^{\frac{1}{p}}.
\end{aligned} \tag{2.8}$$

It has been shown in [7] (see Proposition 2.2 of [7]) that

$$\begin{aligned}
&\sum_{i=1}^{n+1} \int_{\Omega} u_1 x_i \{ (-\Delta)^p (u_1 x_i) - x_i (-\Delta)^p u_1 \} \\
&\leq \left( \lambda_1^{\frac{1}{p}} + n \right)^p - \lambda_1 + 4[2^p - (p+1)] \lambda_1^{\frac{1}{p}} \left( \lambda_1^{\frac{1}{p}} + n \right)^{p-2}.
\end{aligned} \tag{2.9}$$

Inserting (2.8) and (2.9) into (2.7), we infer

$$\begin{aligned}
\sum_{i=1}^{n+1} (\lambda_{i+1} - \lambda_1)^{\frac{1}{2}} \|u_1 \nabla h_i\|^2 &\leq \delta \left\{ \left( \lambda_1^{\frac{1}{p}} + n \right)^p - \lambda_1 + 4[2^p - (p+1)] \lambda_1^{\frac{1}{p}} \left( \lambda_1^{\frac{1}{p}} + n \right)^{p-2} \right\} \\
&\quad + \frac{1}{\delta} \left\{ \lambda_1^{\frac{1}{p}} + \frac{n^2}{4} \right\}.
\end{aligned} \tag{2.10}$$

Minimizing the right hand side of (2.10) as a function of  $\delta$  by choosing

$$\delta = \left( \frac{\lambda_1^{\frac{1}{p}} + \frac{n^2}{4}}{\left( \lambda_1^{\frac{1}{p}} + n \right)^p - \lambda_1 + 4[2^p - (p+1)] \lambda_1^{\frac{1}{p}} \left( \lambda_1^{\frac{1}{p}} + n \right)^{p-2}} \right)^{\frac{1}{2}},$$

we obtain

$$\begin{aligned}
\sum_{i=1}^{n+1} (\lambda_{i+1} - \lambda_1)^{\frac{1}{2}} \|u_1 \nabla h_i\|^2 &\leq \left\{ \left( \lambda_1^{\frac{1}{p}} + n \right)^p - \lambda_1 + 4[2^p - (p+1)] \lambda_1^{\frac{1}{p}} \left( \lambda_1^{\frac{1}{p}} + n \right)^{p-2} \right\}^{\frac{1}{2}} \\
&\quad \times \left\{ 4\lambda_1^{\frac{1}{p}} + n^2 \right\}^{\frac{1}{2}}.
\end{aligned} \tag{2.11}$$

By a transformation of coordinates if necessary, for any point  $q$ , one gets

$$|\nabla h_i|^2 \leq 1 \quad \text{for any } i.$$

It follows that

$$\begin{aligned}
& \sum_{i=1}^{n+1} (\lambda_{i+1} - \lambda_1)^{\frac{1}{2}} |\nabla h_i|^2 \\
&= \sum_{i=1}^n (\lambda_{i+1} - \lambda_1)^{\frac{1}{2}} |\nabla h_i|^2 + (\lambda_{n+1} - \lambda_1)^{\frac{1}{2}} |\nabla h_{n+1}|^2 \\
&= \sum_{i=1}^n (\lambda_{i+1} - \lambda_1)^{\frac{1}{2}} |\nabla h_i|^2 + (\lambda_{n+1} - \lambda_1)^{\frac{1}{2}} \left( n - \sum_{i=1}^n |\nabla h_i|^2 \right) \\
&= \sum_{i=1}^n (\lambda_{i+1} - \lambda_1)^{\frac{1}{2}} |\nabla h_i|^2 + (\lambda_{n+1} - \lambda_1)^{\frac{1}{2}} \sum_{i=1}^n (1 - |\nabla h_i|^2) \\
&\geq \sum_{i=1}^n (\lambda_{i+1} - \lambda_1)^{\frac{1}{2}} |\nabla h_i|^2 + \sum_{i=1}^n (\lambda_{i+1} - \lambda_1)^{\frac{1}{2}} (1 - |\nabla h_i|^2) \\
&= \sum_{i=1}^n (\lambda_{i+1} - \lambda_1)^{\frac{1}{2}}.
\end{aligned} \tag{2.12}$$

From (2.11) and (2.12), we obtain

$$\begin{aligned}
\sum_{i=1}^n (\lambda_{i+1} - \lambda_1)^{\frac{1}{2}} &\leq \left\{ \left( \lambda_1^{\frac{1}{p}} + n \right)^p - \lambda_1 + 4[2^p - (p+1)]\lambda_1^{\frac{1}{p}} \left( \lambda_1^{\frac{1}{p}} + n \right)^{p-2} \right\}^{\frac{1}{2}} \\
&\quad \times \left\{ 4\lambda_1^{\frac{1}{p}} + n^2 \right\}^{\frac{1}{2}},
\end{aligned}$$

which concludes the proof of Theorem.

## References

- [1] Ashbaugh, M.S.: Isoperimetric and universal inequalities for eigenvalues. In spectral theory and geometry (Edinburgh, 1998, E. B. Davies and Yu Safarov, eds.), London Math. Soc. Lecture Notes, 273 (1999), 95-139
- [2] Ashbaugh, M.S., Benguria, R.D.: More bounds on eigenvalue ratios for Dirichlet Laplacians in  $n$  dimensions. SIAM J. Math. Anal. 24, 1622-1651 (1993)
- [3] Brands, J.J.A.M.: Bounds for the ratios of the first three membrane eigenvalues. Arch. Rational Mech. Anal. 16, 265-268 (1964)
- [4] Chen, D.G., Cheng, Q.-M.: Extrinsic estimates for eigenvalues of the Laplace operator. J. Math. Soc. Japan 60, 325-339 (2008)
- [5] Chen, Z.C., Qian, C.L.: Estimates for discrete spectrum of Laplacian operator with any order. J. China Univ. Sci. Tech. 20, 259-266 (1990)
- [6] Cheng, Q.-M., Huang, G.Y., Wei, G.X.: Estimates for lower order eigenvalues of a clamped plate problem. arXiv: 0906.5192

- [7] Cheng, Q.-M., Ichikawa, T., Mametsuka, S.: Estimates for eigenvalues of the poly-Laplacian with any order in a unit sphere. To appear in *Calc. Var.*
- [8] Cheng, Q.-M., Ichikawa, T., Mametsuka, S.: Inequalities for eigenvalues of Laplacian with any order. To appear in *Commun. Contemp. Math.*, 2009
- [9] Huang, G.Y., Li, X.X., Xu, R.W.: Extrinsic estimates for the eigenvalues of Schrödinger operator. To appear in *Geom. Dedicata*
- [10] Payne, L.E., Pólya, G., Weinberger, H.F.: On the ratio of consecutive eigenvalues. *J. Math. Phys.* 35, 289-298 (1956)
- [11] Sun, H.J., Cheng, Q.-M., Yang, H.C.: Lower order eigenvalues of Dirichlet Laplacian. *Manuscripta Math.* 125, 139-156 (2008)